

381,34-3273-9-Rev. 12/04 (PNT 12/04)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID: If **Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If Swallowed:** Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985)878-6751 Monday through Friday 8:00am to 4:30pm CST. After 4:30pm, call your poison control center. For product information call (203)597-1812, Monday - Friday 8:00am to 5:00pm EST. **Note to Physician:** Contains petroleum distillate - vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to any body of water. **PHYSICAL OR CHEMICAL HAZARDS:** Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use in the vicinity of open flame or pilot lights.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product has residual action which keeps on killing for up to 8 weeks after you spray. Remove pets, birds and cover fish aquariums before spraying. Hold container upright. Do not spray up into air. Remove protective cap and direct spray away from face. Hold can 12-15 inches from surface being sprayed. Press button down and spray as directed. Spray surface until slightly wet. **INDOOR: Cockroaches** (including adult and immature stages of both non-resistant and organophosphate and carbamate resistant strains), ants, booklice, boxelder bugs, centipedes, crickets, dermestids, firebrats, fleas, palmetto bugs, silverfish, sowbugs, ticks and waterbugs: Apply thoroughly to all areas where these pests crawl and hide, especially in cracks and crevices, and hidden areas around sinks and storage areas, behind baseboards, around doors and windows, behind and under refrigerators, cabinets, sinks and stoves, the underside of shelves, drawers, bookcases, and similar areas. When pests are driven out, spray directly. Repeat as necessary. **Ants:** Spray freely around ant trails, doors and windows and other places where ants enter the house. **Ticks and Fleas on Surfaces:** Remove pet bedding and destroy or clean thoroughly. Spray pet resting quarters. Adult fleas and larvae contacted by spray will be killed. Put fresh bedding down once spray has dried. Do not spray animals directly. For best results, pets should be treated with an appropriate product before allowing them to return to the treated area. Spray rugs and carpets where infestations are bad. Test first for staining on an inconspicuous piece prior to use. **Bedbugs:** Remove linens and wash before reuse. Apply to mattresses, especially tufts, folds and edges until moist. Apply to the interior of the frame. Allow to dry before remaking bed. Treat baseboards, molding and spot treat floors. **Spiders:** Apply along and behind baseboards, to window and door frames, corners, pipes, storage localities, attics, crawl spaces and other areas over which these pests may crawl. **Centipedes, Ground Beetles, Pillbugs, Sowbugs and Ticks:** Apply around doors and windows and other places where these pests may enter premises. Treat baseboards, storage areas and other locations where these pests are found. **PANTRY PESTS - Grain Beetles (Rusty, Merchant and Saw-toothed), Flour Beetles (Red and Confused), Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Drugstore Beetles, Elmleaf Beetles, Rice Weevils, Lesser Grain Borers, Spider Beetles and Tobacco Moths:** Treat exposed stages. Remove all food-stuffs, utensils and shelf paper from area to be treated. Discard used shelf paper and any infested foodstuffs. Apply spray to shelves and in cracks and crevices behind and under cupboards and cabinets. Allow spray to dry prior to replacement of shelf paper, food and utensils. **Carpet Beetles (Black, Furniture and Varied):** Spray on and under edges of floor coverings, under rugs and furniture and in closets or other localities where these insects are found. **Weaving Clothes Moths:** Use this product as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on clothes. Do not apply to clothing. **Food/Feed Handling Establishments:** For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as a general surface, spot or crack and crevice treatment. This product is authorized for use in Federally Inspected Meat and Poultry Plants as a crack and crevice treatment. Do not use in commercial food/feed handling establishments, restaurants, or other commercial establishments while food is commercially prepared, processed, or served. In the home, all food processing surfaces should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation are also considered food/feed areas. **Non-Food/Non-Feed Areas:** Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be treated. **General Surface Applications:** Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or food/feeds are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food/feed will contact. Clean food/feed handling or processing equipment and thoroughly rinse with clean, fresh water. After application in bakeries, clean all food contact surfaces thoroughly with appropriate equipment. **Spot or Crack & Crevice Applications:** Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made while the facility is in operation provided exposed food is covered or removed from area being treated prior to application. Do not apply directly to food or food-handling surfaces. In the home, do not allow spray to contact food/feed surfaces. If it does, clean surfaces with soap and water. Do not apply directly to food. **Volumetric Treatment of Food Areas and/or Food Handling Establishments:** Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. **OUTDOORS:** Point nozzle away from face. Hold can on a slight downward angle, approximately 12-15 inches from surface to be sprayed. For best results, spray when air is calm. Spray with wind if breeze is blowing. **PESTS ON OUTSIDE SURFACES OF BUILDINGS:** For control of building infestation to **ants, clover mites, crickets and sowbugs (pillbugs).** Spray foundation of building where insects are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include areas such as screens, window frames, eaves, porches, patios, garages and refuse dumps. **CRAWLING INSECTS: Ants, Cockroaches, (including Asian Cockroaches), Centipedes, Crickets, Mole Crickets and Silverfish.** Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel ants. **Ants:** Apply to ant trails, around doors and windows and other places where ants may be found. **Lone Star Ticks, Dog Ticks, Crickets, Fleas and Earwigs:** Apply thoroughly to infestation in bushes, grass or weeds. **Wasps, Mud Daubers, Hornets and Bees:** Spray nests and bees directly. **FLYING INSECTS: Houseflies, Gnats, Mosquitoes and Small Flying Moths:** Use only on outdoor surfaces. Spray outside surfaces of window and door frames and other areas where these pests may enter the home. Also spray localized resting areas, such as under eaves, porches, light fixtures and cords, railings, etc. where these insects may rest. Insects coming to rest on treated surfaces will be killed. **WOOD INFESTING INSECTS: Termites* (Subterranean, Formosan and Drywood)** - For control of above-ground termites in localized areas, apply to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable. Applications may be made to inaccessible areas by drilling, and then injecting with the supplied void injector into the damaged wood or wall voids. These applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement for controlling active infestations of subterranean termites. Termite carton nests in building voids may be injected using a void injector. It is desirable to physically remove carton nest material from building voids when such nests are found. This type of application is intended to be a supplemental treatment for control of drywood and above-ground subterranean termites and other wood-infesting insects. Application to attics, crawl spaces, unfinished basements, or manmade voids may be sprayed to control exposed worker and winged reproductive forms of termites. For control of swarming or reproductive stages of termites, spray to the swarm or area on which they collect. The purpose of such applications of this product is to kill workers or winged reproductive forms which may be present at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection. For control of above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members, drill to find the interior infested cavity and spray using void injector. Termite carton nests in trees may be spray injected with product using the supplied void injector. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from trees when such nests are found. **Carpenter Ants:** For effective control in houses and other structures, apply as general surface, spot, crack and crevice or wall void application. Apply around doors and windows and other places where ants may enter the premises and where they crawl and hide. Spray into cracks and crevices or through small drilled holes into voids where these ants or their nests are present. **For control of carpenter bees** in structures, spray the liquid directly into gallery entrance holes. Following treatment, the entrance holes may be left open 24 hours to be certain that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.

STORAGE AND DISPOSAL

Storage: Do not store near heat or open flame. Store in a cool, dry area away from children. **Container Disposal:** Replace cap, wrap container in several layers of newspaper and discard in trash. Do not incinerate or puncture. Can may be disposed of with other recyclables where local law permits.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



Manufactured by:
Waterbury Companies, Inc.
P.O. Box 1812 Waterbury CT 06722 Made in USA
(800)845-3495 - www.CBProProducts.com



Part No. 34-3273CBB



D-Force™ HPX
RESIDUAL WITH DELTAMETHRIN

CB RESIDUALS

15% VOC Formulation

- Protects Up to 8 Weeks
- Killing Action and Residual Control
- Keeps on Killing For Up to 8 Weeks
- For Indoor/Outdoor Use

Broad Spectrum Insect Control

KILLS: Ants, Bees, Bedbugs, Carpet Beetles, Centipedes, Cockroaches, Crickets, Firebrats, Fleas, Flour Beetles, Grain Beetles, Ground Beetles, Gnats, Midges, Mole Crickets, Moths, Pillbugs, Silverfish, Sowbugs, Spiders, Ticks and Many Other Crawling Insects.

Also Controls and Kills: Wood Infesting Pests - Termites (Subterranean, Formosan & Drywood), Carpenter Ants and Carpenter Bees.

FOR USE IN: Food Handling Establishments, Including Food Service, Manufacturing and Processing Establishments, Such as Restaurants, Cafeterias, Supermarkets, Bakeries, Breweries, Dairies, Meat, Poultry & Egg Processing & Packaging Plants, Canneries, Feed Handling Establishments Including Feed Manufacturing and Processing Establishments.

Also For Use in: Homes, Tents, Aircraft, Apartments, Atriums, Attics, Automobiles, Basements, Bathrooms, Boats, Bottling facilities, Buses, Cabins, Campers, Candy Plants, Carports, Cat Sleeping Quarters, Clothes Storage, Condominiums, Decks, Dens, Driveways, Garages, Gazebos, Granaries & Grain Mills, Hospitals, Hotels, Industrial Buildings, Kitchens, Laboratories, Lanais, Living Rooms, Mausoleums, Offices, Parlors, Patios, Pet Sleeping Areas, Play Rooms, Porches, Railcars, Recreational Vehicles, Rooms, Schools, Ships & Vessels, Solariums, Storage Areas, Sun Rooms, Trailers, Trucks, Utility Rooms, Verandahs, Warehouses and Wineries.

ACTIVE INGREDIENT:

Deltamethrin.....	0.06%
INERT INGREDIENTS.....	99.94%
	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional
Precautionary Statements.

EPA Reg No. 9444- 217 EPA Est. No. 9444-LA-1
U.S. patents pending.

NET WEIGHT 14 oz. / 397 g

